

Facts about Anxiety Disorders

Anxiety Disorders affect about 40 million American adults age 18 years and older (about 18%) in a given year, causing them to be filled with fearfulness and uncertainty. Unlike the relatively mild, brief anxiety caused by a stressful event (such as speaking in public or a first date), anxiety disorders last at least 6 months and can get worse if they are not treated. Anxiety disorders commonly occur along with other mental or physical illnesses, including alcohol or substance abuse, which may mask anxiety symptoms or make them worse. In some cases, these other illnesses need to be treated before a person will respond to treatment for the anxiety disorder.

Effective therapies for anxiety disorders are available, and research is uncovering new treatments that can help most people with anxiety disorders lead productive, fulfilling lives. Each anxiety disorder has different symptoms, but all the symptoms cluster around excessive, irrational fear and dread.

People with **generalized anxiety disorder (GAD)** go through the day filled with exaggerated worry and tension, even though there is little or nothing to provoke it. They anticipate disaster and are overly concerned about health issues, money, family problems, or difficulties at work. Sometimes just the thought of getting through the day produces anxiety.

GAD is diagnosed when a person worries excessively about a variety of everyday problems for at least 6 months. People with GAD can't seem to get rid of their concerns, even though they usually realize that their anxiety is more intense than the situation warrants. They can't relax, startle easily, and have difficulty concentrating. Often they have trouble falling asleep or staying asleep. Physical symptoms that often accompany the anxiety include fatigue, headaches, muscle tension, muscle aches, difficulty swallowing, trembling, twitching, irritability, sweating, nausea, lightheadedness, having to go to the bathroom frequently, feeling out of breath, and hot flashes.

When their anxiety level is mild, people with GAD can function socially and hold down a job. Although they don't avoid certain situations as a result of their disorder, people with GAD can have difficulty carrying out the simplest daily activities if their anxiety is severe.

GAD affects about 6.8 million American adults, including twice as many women as men. The disorder develops gradually and can begin at any point in the life cycle, although the years of highest risk are between childhood and middle age. There is evidence that genes play a modest role in GAD.

Other anxiety disorders, depression, or substance abuse often accompany GAD, which rarely occurs alone. GAD is commonly treated with medication or cognitive-behavioral therapy, but co-occurring conditions must also be treated using the appropriate therapies.

If you or a family member is experiencing a mental health or an alcohol or other drug-related emergency, seek immediate assistance by calling the 24-hour Suicide Prevention, Mental Health Crisis, Information and Referral Hotline: (216) 623-6888 or the United Way's First Call for Help, 211 or (216) 436-2000.